




How to Use AzaSite®

Before using AzaSite®, thoroughly wash your hands. Then follow the steps listed below.

Please follow steps 1 and 2 the *first* time you use a new bottle of AzaSite®:



1. Remove the white tamper-evident cap by turning it in a clockwise direction. Discard the white cap.




2. Holding the bottle upright, remove the tan cap. This will allow any excess pressure in the bottle to be released. Replace the tan cap on the bottle and close securely.

Please follow steps 3 through 6 *each* time you use AzaSite®:



3. Turn the closed bottle upside down.



4. Shake hand firmly before each use to move the contents into the tip of the bottle.



5. Keeping the bottle upside down, remove the tan cap.



6. Tilt head back. With the bottle inverted, gently squeeze the bottle to instill 1 drop into each affected eye. Replace the tan cap back on the bottle and close securely after each administration for proper storage.

Just remember—**Flip, Whip, and Drip.**

AzaSITE®
(azithromycin ophthalmic solution) 1%

What side effects might I experience with AzaSite®?

In clinical trials, the most common adverse event was eye irritation, which occurred in 1% to 2% of patients. Other adverse events (occurring in <1% of patients) were burning, stinging, and eye irritation upon instillation; skin irritation or rash; corneal erosion; dry eye; altered taste; nasal congestion; eye discharge; inflammation of the cornea; and sinus infection.

What if I have questions about AzaSite®?

Contact your doctor if you have any questions about AzaSite®. You can also visit www.azasite.com, or you may call **1-888-881-4696** to speak with a healthcare representative.

AzaSITE[®]
(azithromycin ophthalmic solution) 1%

AzaSite® (azithromycin ophthalmic solution) 1% is indicated for the treatment of bacterial conjunctivitis (pinkeye) in adults and children 1 year of age and older. Recommended dosing for AzaSite® is 1 drop twice a day for 2 days, followed by 1 drop once a day for the next 5 days, for a total of 9 drops per affected eye.

Important Safety Information: For use as an eyedrop only.

Please see accompanying full Prescribing Information.

INSPIRE 

©2008 AZA-0257 May 2008 Inspire Pharmaceuticals, Inc.
AzaSite is a registered trademark of InSite Vision, Inc. All rights reserved.